arXiv:1610.09783v1 [math.CO] 31 Oct 2016

Hermitian-Randić matrix and Hermitian-Randić energy of mixed graphs *

Yong Lu[1], Ligong Wang[2,†]and Qiannan Zhou[3]
Department of Applied Mathematics, School of Science, Northwestern Polytechnical University, Xi'an, Shaanxi 710072, People's Republic of China.
[1] E-mail: luyong.gougou@163.com
[2] E-mail: lgwangmath@163.com
[3] E-mail: qnzhoumath@163.com

Abstract

Let M be a mixed graph and $H(M)$ be its Hermitian-adjacency matrix. If we add every edge and arc in M a Randić weight, then we can get a new weighted Hermitian-adjacency matrix. What are the properties of this new matrix? Motivated by this, we define the Hermitian-Randić matrix $R_H(M) = (r_h)_{kl}$ of a mixed graph M, where $(r_h)_{kl} = -(r_h)_{lk} = \frac{\mathbf{i}}{\sqrt{d_k d_l}}$ ($\mathbf{i} = \sqrt{-1}$) if (v_k, v_l) is an arc of M, $(r_h)_{kl} = (r_h)_{lk} = \frac{1}{\sqrt{d_k d_l}}$ if $v_k v_l$ is an undirected edge of M, and $(r_h)_{kl} = 0$ otherwise. In this paper, firstly, we compute the characteristic polynomial of the Hermitian-Randić matrix of a mixed graph. Furthermore, we give bounds to the Hermitian-Randić energy of a general mixed graph. Finally, we give some results about the Hermitian-Randić energy of mixed trees.

Key Words: Mixed graph, Hermitian-adjacency matrix, Hermitian-Randić matrix, Hermitian-Randić energy.

AMS Subject Classification (2010): 05C50, 05C07, 05C31.

1 Introduction

In this paper, we only consider simple graphs without multiedges and loops. A graph M is said to be *mixed* if it is obtained from an undirected graph M_U by orienting a subset of its edges. We call M_U the *underlying graph* of M. Clearly, a mixed graph concludes both possibilities of all edges oriented and all edges undirected as extreme cases.

Let M be a mixed graph with vertex set $V(M) = \{v_1, v_2, \ldots, v_n\}$ and edge set $E(M)$. For $v_i, v_j \in V(M)$, we denote an undirected edge joining two vertices v_i and v_j of M by $v_i v_j$ (or $v_i \leftrightarrow v_j$). Denote a directed edge (or arc) from v_i to v_j by (v_i, v_j) (or $v_i \to v_j$). In addition, let $E_0(M)$ denote the set of all undirected edges and $E_1(M)$ denote all the directed arcs set. Clearly, $E(M)$ is the union of $E_0(M)$ and $E_1(M)$. A mixed graph is called *mixed tree* (or *mixed bipartite graph*) if its underlying graph is a tree (or bipartite graph). In general, the

*Supported by the National Natural Science Foundation of China (No. 11171273)
†Corresponding author.

order, size, number of components, degree of a vertex of M are the same to those in M_U. We use Bondy and Murty [2] for terminologies and notations not defined here.

Let G be a simple graph with vertex set $\{v_1, v_2, \ldots, v_n\}$. The *adjacency matrix* of a simple graph G of order n is defined as the $n \times n$ symmetric square matrix $A = A(G) = (a_{ij})$, where $a_{ij} = 1$ if v_iv_j is an edge of G, otherwise $a_{ij} = 0$. We denote by $d_i = d(v_i) = d_G(v_i)$ $(i = 1, 2, \ldots, n)$ the degree of vertex v_i. In addition, for a mixed graph M, if $v_i \in V(M)$, then we also denote by $d_i = d(v_i) = d_{M_U}(v_i)$. The *energy* of the graph G (See the survey of Gutman, Li and Zhang [15] and the book of Li, Shi and Gutman [21]) is defined as $\mathcal{E}_A(G) = \sum_{i=1}^{n} |\rho_i|$, where $\rho_1, \rho_2, \ldots, \rho_n$ are all eigenvalues of $A(G)$.

A convenient parameter of G is the *general Randić index* $R_\alpha(G)$ defined as $R_\alpha(G) = \sum_{uv \in E(G)} (d_u d_v)^\alpha$, where the summation is over all (unordered) edges uv in G. The molecular structure-descriptor, first proposed by Randić [27] in 1975, is defined as the sum of $\frac{1}{\sqrt{d_u d_v}}$ over all edges uv of G (with $\alpha = -\frac{1}{2}$). Nowadays, $R = R(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_u d_v}}$ of G is referred to as the *Randić index*. Countless chemical applications, the mathematical properties and mathematical chemistry of the Randić index were reported in [18, 20, 28].

Gutman et al.[14] pointed out that the Randić-index-concept is purposeful to associate the graph G a symmetric square matrix of order n, named *Randić matrix* $R(G) = (r_{ij})$, where $r_{ij} = \frac{1}{\sqrt{d_i d_j}}$ if v_iv_j is an edge of G, otherwise $r_{ij} = 0$. Let $D(G)$ be the diagonal matrix of vertex degrees of G. If G has no isolated vertices, then $R(G) = D(G)^{-\frac{1}{2}} A(G) D(G)^{-\frac{1}{2}}$.

The concept of the *Randić energy* of a graph G, denoted by $\mathcal{E}_R(G)$, was introduced in [6] as $\mathcal{E}_R(G) = \sum_{i=1}^{n} |\gamma_i|$, where γ_i is the eigenvalues of $R(G)$, $i = 1, 2, \ldots, n$. Some basic properties of the Randić index, Randić matrix and Randić energy were determined in the papers [3, 4, 5, 6, 10, 11, 12, 14, 17, 22, 23, 24, 29].

An oriented graph G^σ is a digraph which assigns each edge of G with a direction σ. The *skew adjacency matrix* associated to G^σ is the $n \times n$ matrix $S(G^\sigma) = (s_{ij})$, where $s_{ij} = -s_{ji} = 1$ if (v_i, v_j) is an arc of G^σ, otherwise $s_{ij} = s_{ji} = 0$. The *skew energy* of G^σ, denoted by $\mathcal{E}_S(G^\sigma)$, is defined as the sum of the norms of all the eigenvalues of $S(G^\sigma)$. For more details about skew energy we can refer to [1, 19].

In 2016, Gu, Huang and Li [11] defined the *skew Randić matrix* $R_s(G^\sigma) = ((r_s)_{ij})$ of an oriented graph G^σ of order n, where $(r_s)_{ij} = -(r_s)_{ji} = \frac{1}{\sqrt{d_i d_j}}$ if (v_i, v_j) is an arc of G^σ, otherwise $(r_s)_{ij} = (r_s)_{ji} = 0$. Let $D(G)$ be the diagonal matrix of vertex degrees of G. If G^σ has no isolated vertices, then $R_s(G^\sigma) = D(G)^{-\frac{1}{2}} S(G^\sigma) D(G)^{-\frac{1}{2}}$.

The *Hermitian-adjacency matrix* of a mixed graph M of order n is the $n \times n$ matrix $H(M) = (h_{kl})$, where $h_{kl} = -h_{lk} = \mathbf{i}$ $(\mathbf{i} = \sqrt{-1})$ if (v_k, v_l) is an arc of M, $h_{kl} = h_{lk} = 1$ if v_kv_l is an undirected edge of M, and $h_{kl} = 0$ otherwise. Obviously, $H(M) = H(M)^* := \overline{H(M)}^T$. Thus all its eigenvalues are real. This matrix was introduced by Liu and Li in [25] and independently by Guo and Mohar in [13]. The *Hermitian energy* of a mixed graph M is defined as $\mathcal{E}_H(M) = \sum_{i=1}^{n} |\lambda_i|$, where $\lambda_1, \lambda_2, \ldots, \lambda_n$ are all eigenvalues of $H(M)$. Denote by $Sp_H(M) = (\lambda_1, \lambda_2, \ldots, \lambda_n)$ the *spectrum* of $H(M)$. For more details about Hermitian-adjacency matrix and Hermitian energy of mixed graphs, we can refer to [7, 13, 25, 26, 30, 31].

From the above we can see that if we add every edge in a simple graph G a Randić weight, then we can get a Randić matrix $R(G)$. If we add every arc in an oriented graph G^σ a Randić weight, then we can get a skew Randić matrix $R_s(G^\sigma)$. Let M be a mixed graph and $H(M)$ be its Hermitian-adjacency matrix, if we add every edge and arc in M a Randić weight, then we can get a new weighted Hermitian-adjacency matrix. What are the properties of this new

matrix? Motivated by this, we define the Hermitian-Randić matrix of a mixed graph M.

Let M be a mixed graph on the vertex set $\{v_1, v_2, \ldots, v_n\}$, then the *Hermitian-Randić matrix* of M is the $n \times n$ matrix $R_H(M) = ((r_h)_{kl})$, where

$$(r_h)_{kl} = \begin{cases} \frac{1}{\sqrt{d_k d_l}}, & \text{if } v_k \leftrightarrow v_l, \\ \frac{\mathbf{i}}{\sqrt{d_k d_l}}, & \text{if } v_k \rightarrow v_l, \\ \frac{-\mathbf{i}}{\sqrt{d_k d_l}}, & \text{if } v_l \rightarrow v_k, \\ 0, & \text{otherwise.} \end{cases}$$

Let $D(M_U)$ be the diagonal matrix of vertex degrees of M_U. If M has no isolated vertices, then $R_H(M) = D(M_U)^{-\frac{1}{2}} H(M) D(M_U)^{-\frac{1}{2}}$. For a mixed graph M, let $R_H(M)$ be its Hermitian-Randić matrix. It is obvious that $R_H(M)$ is Hermitian matrix, so all its eigenvalues $\mu_1, \mu_2, \ldots, \mu_n$ are real. The *spectrum* of $R_H(M)$ is defined as $Sp_{R_H}(M) = (\mu_1, \mu_2, \ldots, \mu_n)$. The energy of $R_H(M)$, denoted by $\mathcal{E}_{R_H}(M)$, called *Hermitian-Randić energy*, which is defined as the sum of the absolute values of its eigenvalues of $R_H(M)$, that is $\mathcal{E}_{R_H}(M) = \sum_{i=1}^{n} |\mu_i|$.

2 Hermitian-Randić characteristic polynomial of a mixed graph

In this section, we will give the characteristic polynomial of Hermitian-Randić matrix of a mixed graph M, i.e., the R_H-characteristic polynomial of M. At first, we will introduce some basic definitions.

The *value* of a mixed walk $W = v_1 v_2 \cdots v_l$ is $r_h(W) = (r_h)_{12}(r_h)_{23}(r_h)_{(l-1)l}$. A mixed walk W is *positive* (or *negative*) if $r_h(W) = \frac{1}{\sqrt{d_1 d_l} d_2 d_3 \cdots d_{(l-1)}}$ (or $r_h(W) = -\frac{1}{\sqrt{d_1 d_l} d_2 d_3 \cdots d_{(l-1)}}$). Note that for one direction the value of a mixed walk or a mixed cycle is α, then for the reversed direction its value is $\overline{\alpha}$. Thus, if the value of a mixed cycle C is $\prod_{v_j \in V(C)} \frac{1}{d(v_j)}$ (resp. $-\prod_{v_j \in V(C)} \frac{1}{d(v_j)}$) in a direction, then its value is $\prod_{v_j \in V(C)} \frac{1}{d(v_j)}$ (resp. $-\prod_{v_j \in V(C)} \frac{1}{d(v_j)}$) for the reversed direction. In these situations, we just term this mixed cycle as a positive (resp. negative) mixed cycle without mentioning any direction.

If each mixed cycle is positive (resp. negative) in a mixed graph M, then M is *positive* (resp. *negative*). A mixed graph M is called an *elementary graph* if every component of M is an edge, an arc or a mixed cycle, where every edge-component in M is defined to be positive. A *real spanning elementary subgraph* of a mixed graph M is an elementary subgraph such that it contains all vertices of M and all its mixed cycles are real.

Now we will give two results which are similar to those of [9, 16, 25].

Let M be a mixed graph of order n with its Hermitian-Randić matrix $R_H(M)$. Denote the *R_H-characteristic polynomial* of $R_H(M)$ of M by

$$P_{R_H}(M, x) = \det(xI - R_H(M)) = x^n + a_1 x^{n-1} + a_2 x^{n-2} + \cdots + a_n.$$

Theorem 2.1. *Let $R_H(M)$ be the Hermitian-Randić matrix of a mixed graph M of order n. Then*

$$\det R_H(M) = \sum_{M'} (-1)^{r(M')+l(M')} 2^{s(M')} W(M'),$$

where the summation is over all real spanning elementary subgraphs M' of M, $r(M') = n - c(M')$, $c(M')$ denotes the number of components of M', $l(M')$ denotes the number of negative mixed cycles of M', $s(M')$ denotes the number of mixed cycles with length ≥ 3 in M', $W(M') = \prod_{v_i \in V(M')} \frac{1}{d_{M_U}(v_i)}$.

Proof. Let M be a mixed graph of order n with vertex set $\{v_1, v_2, \ldots, v_n\}$. Then

$$\det R_H(M) = \sum_{\pi \in S_n} sgn(\pi)(r_h)_{1\pi(1)}(r_h)_{2\pi(2)} \cdots (r_h)_{n\pi(n)},$$

where S_n is the set of all permutations on $\{1, 2, \ldots, n\}$.

Consider a term $sgn(\pi)(r_h)_{1\pi(1)}(r_h)_{2\pi(2)} \cdots (r_h)_{n\pi(n)}$ in the expansion of $\det R_H(M)$. If $v_k v_{\pi(k)}$ is not an edge or arc of M, then $(r_h)_{k\pi(k)} = 0$; that is, this term vanishes. Thus, if the term corresponding to a permutation π is non-zero, then π is fixed-point-free and can be expressed uniquely as the composition of disjoint cycles of length at least 2. Consequently, each non-vanishing term in the expansion of $\det R_H(M)$ gives rise to an elementary mixed graph M' of M, with $V(M') = V(M)$. That is, M' is a spanning elementary subgraph of M of order n.

A spanning elementary subgraph M' of M with $s(M')$ number of mixed cycles (length ≥ 3) gives $2^{s(M')}$ permutations π, since for each mixed cycle-component in M' there are two ways of choosing the corresponding cycles in π. For a vertex $v_k \in V(M')$, we denote by $d_k = d(v_k) = d_{M_U}(v_k)$. Furthermore, if for some direction of a permutation π, a mixed cycle-component C_1 has value $\mathbf{i}\prod_{v_j \in V(C_1)} \frac{1}{d(v_j)}$ (or $-\mathbf{i}\prod_{v_j \in V(C_1)} \frac{1}{d(v_j)}$), then for the other direction C_1 has value $-\mathbf{i}\prod_{v_j \in V(C_1)} \frac{1}{d(v_j)}$ (or $\mathbf{i}\prod_{v_j \in V(C_1)} \frac{1}{d(v_j)}$) and vice verse. Thus, they cancel each other in the summation. In addition, if for some direction of a permutation π, C_1 has value $\prod_{v_j \in V(C_1)} \frac{1}{d(v_j)}$ (or $-\prod_{v_j \in V(C_1)} \frac{1}{d(v_j)}$), then for the other direction C_1 has the same value. For each edge-component (kl) corresponds to the factors $(r_h)_{kl}(r_h)_{lk}$ has value $\frac{1}{\sqrt{d_k d_l}}\frac{1}{\sqrt{d_l d_k}} = \frac{1}{d_k d_l}$. For each arc-component (kl) corresponds to the factors $(r_h)_{kl}(r_h)_{lk}$ has value $\frac{\mathbf{i}\cdot(-\mathbf{i})}{\sqrt{d_k d_l}\sqrt{d_l d_k}} = \frac{1}{d_k d_l}$.

Since $sgn(\pi) = (-1)^{n-c(M')} = (-1)^{r(M')}$ and each real spanning elementary subgraph M' contributes $(-1)^{r(M')+l(M')}2^{s(M')}\prod_{v_i \in V(M')} \frac{1}{d_{M_U}(v_i)}$ to the determinant of $R_H(M)$. This completes the proof. ■

Now, we shall obtain a description of all the coefficients of the characteristic polynomial $P_{R_H}(M, x)$ of a mixed graph M.

Theorem 2.2. *For a mixed graph M, if the R_H-characteristic polynomial of M is denoted by $P_{R_H}(M, x) = \det(xI - R_H(M)) = x^n + a_1 x^{n-1} + a_2 x^{n-2} + \cdots + a_n$, then the coefficients of $P_{R_H}(M, x)$ are given by*

$$(-1)^k a_k = \sum_{M'} (-1)^{r(M')+l(M')} 2^{s(M')} \prod_{v_i \in V(M')} \frac{1}{d_{M_U}(v_i)},$$

where the summation is over all real elementary subgraphs M' with order k of M, $r(M') = k - c(M')$, $c(M')$ denotes the number of components of M', $l(M')$ denotes the number of negative mixed cycles of M', $s(M')$ denotes the number of mixed cycles with length ≥ 3 in M'.

Proof. The proof follows from Theorem 2.1 and the fact that $(-1)^k a_k$ is the summation of determinants of all principal $k \times k$ submatrices of $R_H(M)$. ■

From Theorem 2.2, we can obtain the following results.

Corollary 2.3. *For a mixed graph M, let the R_H-characteristic polynomial of M be denoted by $P_{R_H}(M, x) = |xI - R_H(M)| = x^n + a_1 x^{n-1} + a_2 x^{n-2} + \cdots + a_n$.*

(1) If M is a mixed tree, then

$$(-1)^k a_k = \sum_{M'} (-1)^{r(M')} \prod_{v_i \in V(M')} \frac{1}{d_{M_U}(v_i)}.$$

(2) If M is a mixed graph and its underlying graph M_U is r regular ($r \neq 0$), then

$$(-1)^k a_k = \sum_{M'} (-1)^{r(M')+l(M')} 2^{s(M')} \frac{1}{r^k}.$$

(3) If M is a mixed bipartite graph, then all coefficients of a_{odd} are equal to 0, and its spectrum is symmetry about 0.

Note that if M is a positive mixed graph, then for every real elementary subgraph M' of M, we have

$$\begin{aligned}
&(-1)^{r(M')+l(M')} 2^{s(M')} \prod_{v_i \in V(M')} \frac{1}{d_{M_U}(v_i)} \\
&= (-1)^{r(M')} 2^{s(M')} \prod_{v_i \in V(M')} \frac{1}{d_{M_U}(v_i)} \\
&= (-1)^{r(M'_U)} 2^{s(M'_U)} \prod_{v_i \in V(M'_U)} \frac{1}{d_{M_U}(v_i)}.
\end{aligned}$$

Then $P_{R_H}(M, x) = P_{R_H}(M_U, x)$, that is to say

Theorem 2.4. *If M is a positive mixed graph and M_U be its underlying graph, then $Sp_{R_H}(M) = Sp_{R_H}(M_U)$.*

3 Bounds on Hermitian-Randić energy of mixed graphs

In this section, we will give some bounds about Hermitian-Randić energy of mixed graphs. First, we will give some properties of Hermitian-Randić matrix of mixed graphs.

Lemma 3.1. *Let M be a mixed graph of order $n \geq 1$.*

(1) $\mathcal{E}_{R_H}(M) = 0$ if and only if $M \cong \overline{K}_n$.

(2) If $M = M_1 \cup M_2 \cup \cdots \cup M_p$, then $\mathcal{E}_{R_H}(M) = \mathcal{E}_{R_H}(M_1) + \mathcal{E}_{R_H}(M_2) + \cdots + \mathcal{E}_{R_H}(M_p)$.

From Lemma 3.1, we can obtain the following theorem.

Theorem 3.2. *Let M be a mixed graph with vertex set $V(M) = \{v_1, v_2, \ldots, v_n\}$ and d_k is the degree of v_k, $k = 1, 2, \ldots, n$. Let $H(M)$ and $R_H(M)$ be the Hermitian-adjacency matrix and Hermitian-Randić matrix of M, respectively. If M has isolated vertices, then* $\det H(M) = \det R_H(M) = 0$. *If M has no isolated vertices, then*

$$\det R_H(M) = \frac{1}{d_1 d_2 \cdots d_n} \det H(M).$$

Proof. If M has l isolated vertices, then $M = M' \cup \overline{K}_l$, where M' has no isolated vertices. By Lemma 3.1, we have $Sp_{R_H}(M) = Sp_{R_H}(M') \cup \{0, l\ times\}$ and an analogous relation holds for Hermitian-adjacency spectrum of M. That is, $H(M)$ and $R_H(M)$ have zero eigenvalues, therefore their determinants are equal to zero. If M has no isolated vertices, then $R_H(M) = D(M_U)^{-\frac{1}{2}} H(M) D(M_U)^{-\frac{1}{2}}$ is applicable, where $D(M_U)$ is the diagonal matrix of vertex degrees. The matrices $R_H(M)$ and $D(M_U)^{-\frac{1}{2}} R_H(M) D(M_U)^{\frac{1}{2}}$ are similar and thus have equal eigenvalues. We have

$$D(M_U)^{-\frac{1}{2}} R_H(M) D(M_U)^{\frac{1}{2}} = D(M_U)^{-1} H(M),$$

and therefore,

$$\det R_H(M) = \det[D(M_U)^{-1} H(M)] = \det D(M_U)^{-1} \det H(M).$$

So,

$$\det R_H(M) = \frac{1}{d_1 d_2 \cdots d_n} \det H(M).$$

This completes the proof. ■

Similar to Theorem 3.2, we can obtain the following theorem.

Theorem 3.3. *If M is a mixed graph with vertex set $V(M) = \{v_1, v_2, \ldots, v_n\}$ and its underlying graph M_U is r regular, then $\mathcal{E}_{R_H}(M) = \frac{1}{r}\mathcal{E}_H(M)$. In addition, if $r = 0$, then $\mathcal{E}_{R_H}(M) = 0$.*

Proof. If $r = 0$, then M is the graph that has no edges. Then all the entries of $R_H(M)$ are equal to 0, i.e., $R_H(M) = \boldsymbol{0}$. Similarly, $H(M) = \boldsymbol{0}$. Since all eigenvalues of the zero matrix are equal to 0. Hence, $\mathcal{E}_{R_H}(M) = \mathcal{E}_H(M)$=0.

If $r > 0$, i.e., M is regular of degree $r > 0$. Then $d_1 = d_2 = \cdots = d_n = r$, where d_k is the degree of v_k, $k = 1, 2, \ldots, n$. Hence

$$(r_h)_{sk} = \begin{cases} \frac{1}{r}, & v_s \leftrightarrow v_k, \\ \frac{\mathbf{i}}{r}, & v_s \rightarrow v_k, \\ \frac{-\mathbf{i}}{r}, & v_k \rightarrow v_s, \\ 0, & \text{otherwise.} \end{cases}$$

Which implies that $R_H(M) = \frac{1}{r} H(M)$. Therefore, $\mu_i = \frac{1}{r}\lambda_i$, where μ_i is the eigenvalue of $R_H(M)$, and λ_i is the eigenvalue of $H(M)$, for $i = 1, 2, \ldots, n$. Then this theorem follows from the definitions of $\mathcal{E}_{R_H}(M)$ and $\mathcal{E}_H(M)$. ■

Similar the results about the skew Randić energy in [11], we can establish the following lower and upper bounds for the Hermitian-Randić energy. First, we need the following theorem. Here and later by $\mathbf{I}_n$ is denoted by the unit matrix of order n.

Theorem 3.4. *Let M be a mixed graph of order n and $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_n$ be the Hermitian-Randić spectrum of $R_H(M)$. Then $|\mu_1| = |\mu_2| = \cdots = |\mu_n|$ if and only if there exists a constant $c = |\mu_i|^2$ for all i such that $R_H^2(M) = c\mathbf{I}_n$.*

Proof. Let $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_n$ be the Hermitian-Randić spectrum of $R_H(M)$. Then there exists a unitary matrix U such that

$$U^*R_H(M)U = U^*R_H(M)^*U = \begin{pmatrix} \mu_1 & & \\ & \ddots & \\ & & \mu_n \end{pmatrix}.$$

So,

$$\begin{aligned} & |\mu_1| = |\mu_2| = \cdots = |\mu_n| \\ \Leftrightarrow\ & U^*R_H(M)^*R_H(M)U = c\mathbf{I}_n \\ \Leftrightarrow\ & U(U^*R_H(M)^*R_H(M)U)U^* = cUU^* \\ \Leftrightarrow\ & R_H(M)^*R_H(M) = c\mathbf{I}_n \\ \Leftrightarrow\ & R_H^2(M) = c\mathbf{I}_n, \end{aligned}$$

where c is a constant and $c = |\mu_i|^2$ for all i.

This completes the proof. ■

Theorem 3.5. *Let M be a mixed graph of order n and $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_n$ be the Hermitian-Randić spectrum of $R_H(M)$. Let M_U be the underlying graph of M, $p = |\det R_H(M)|$. Then*

$$\sqrt{2R_{-1}(M_U) + n(n-1)p^{\frac{2}{n}}} \leq \mathcal{E}_{R_H}(M) \leq \sqrt{2nR_{-1}(M_U)},$$

with equalities hold both in the lower bound and upper bound if and only if there exists a constant $c = |\mu_i|^2$ for all i such that $R_H^2(M) = c\mathbf{I}_n$.

Proof. Let $\{\mu_1, \mu_2, \ldots, \mu_n\}$ be the Hermitian-Randić spectrum of M, where $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_n$. Since $\sum_{j=1}^n \mu_j^2 = tr(R_H^2(M)) = \sum_{j=1}^n \sum_{k=1}^n (r_h)_{jk}(r_h)_{kj} = \sum_{j=1}^n \sum_{k=1}^n (r_h)_{jk}\overline{(r_h)}_{jk}$ $= \sum_{j=1}^n \sum_{k=1}^n |(r_h)_{jk}|^2 = 2R_{-1}(M_U)$, where $R_{-1}(M_U) = \sum_{v_jv_k \in E(M_U)} \frac{1}{d_jd_k}$ (unordered).

Applying the Cauchy-Schwartz inequality we have

$$\mathcal{E}_{R_H}(M) = \sum_{j=1}^n |\mu_j| \leq \sqrt{\sum_{j=1}^n |\mu_j|^2} \cdot \sqrt{n} = \sqrt{2nR_{-1}(M_U)}.$$

On the other hand,

$$|\mathcal{E}_{R_H}(M)|^2 = \left(\sum_{j=1}^n |\mu_j|\right)^2 = \sum_{j=1}^n |\mu_j|^2 + \sum_{1 \leq i \neq j \leq n} |\mu_i||\mu_j|.$$

By using arithmetic geometric average inequality, we can get that

$$|\mathcal{E}_{R_H}(M)|^2 = \sum_{j=1}^n |\mu_j|^2 + \sum_{1 \leq i \neq j \leq n} |\mu_i||\mu_j| \geq 2R_{-1}(M_U) + n(n-1)p^{\frac{2}{n}}.$$

Therefore, we can obtain the lower bound on Hermitian-Randić energy,

$$\mathcal{E}_{R_H}(M) \geq \sqrt{2R_{-1}(M_U) + n(n-1)p^{\frac{2}{n}}}.$$

From the Cauchy-Schwartz inequality and arithmetic geometric average inequality, we know that the equalities hold both in the lower bound and upper bound if and only if $|\mu_1| = |\mu_2| = \cdots = |\mu_n|$, i.e., there exists a constant $c = |\mu_i|^2$ for all i such that $R_H^2(M) = c\mathbf{I}_n$.

This completes the proof. ■

Corollary 3.6. *Let M be a mixed graph and its underlying graph M_U be r ($\neq 0$) regular and $E(M_U) = m$. Let $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_n$ be the Hermitian-Randić spectrum of $R_H(M)$. Then*

$$\sqrt{\frac{n}{r} + n(n-1)p^{\frac{2}{n}}} \leq \mathcal{E}_{R_H}(M) \leq \frac{n\sqrt{r}}{r},$$

where $p = |\det R_H(M)|$, with equalities hold both in the lower bound and upper bound if and only if $\frac{1}{r} = |\mu_i|^2$ for all i such that $R_H^2(M) = \frac{1}{r}\mathbf{I}_n$.

Proof. If M is a mixed graph and its underlying graph M_U is r regular, then $R_{-1}(M_U) = \frac{m}{r^2}$ and $2m = nr$. By Theorems 3.4 and 3.5, we can obtain the results. ■

Lemma 3.7. *[24] Let G be a graph of order n with no isolated vertices. Then*

$$\frac{n}{2(n-1)} \leq R_{-1}(G) \leq \lfloor \frac{n}{2} \rfloor,$$

with equality in the lower bound if and only if G is a complete graph, and equality in the upper bound if and only if either

(1) n is even and G is the disjoint union of $n/2$ paths of length 1, or

(2) n is odd and G is the disjoint union of $(n-3)/2$ paths of length 1 and one path of length 2.

Combining with Theorem 3.5 and Lemma 3.7, we can get upper and lower bounds for the Hermitian-Randić energy by replacing $R_{-1}(M_U)$ with other parameters. We now give bounds of Hermitian-Randić energy of a mixed graph with respect to its order.

Theorem 3.8. *Let M be a mixed graph of order $n \geq 3$ without isolated vertices and M_U be its underlying graph. Let $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_n$ be the Hermitian-Randić spectrum of $R_H(M)$. Then*

$$\sqrt{\frac{2n}{n-1}} \leq \mathcal{E}_{R_H}(M) \leq n.$$

The equality in the upper bound holds if and only if n is even and M_U is the disjoint union of $n/2$ paths of length 1. The equality in the lower bound holds if and only if M_U is a complete graph and $\mu_1 = -\mu_n \neq 0$, $\mu_j = 0$, $j = 2, \ldots, n-1$.

Proof. Let $R_H(M)$ be the Hermitian-Randić matrix of M and $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_n$ be the Hermitian-Randić spectrum of $R_H(M)$.

For the upper bound, combining with Lemma 3.7 and $\mathcal{E}_{R_H}(M) \leq \sqrt{2nR_{-1}(M_U)}$ of Theorem 3.5, we have

$$\mathcal{E}_{R_H}(M) \leq \sqrt{2nR_{-1}(M_U)} \leq \sqrt{2n\lfloor \frac{n}{2} \rfloor} \leq n.$$

From Theorem 3.5 and Lemma 3.7, we know that the equality in the upper bound holds if and only if n is even, M_U is the graph described in Lemma 3.7 (1), and $|\mu_1| = |\mu_2| = \cdots = |\mu_n|$,

that is, we can obtain the upper bound when n is even and M_U is the disjoint union of $n/2$ paths of length 1.

For the lower bound, since the sum of the diagonal entries of $R_H(M)$ is 0, i.e., $\sum_{k=1}^{n} \mu_k = 0$. Then

$$\begin{aligned}\left(\sum_{k=1}^{n} \mu_k\right)\left(\sum_{l=1}^{n} \mu_l\right) &= \sum_{k=1}^{n} \mu_k^2 + \sum_{1\le k\ne l\le n} \mu_k\mu_l \\ &= \sum_{k=1}^{n} \mu_k^2 + 2\sum_{k<l} \mu_k\mu_l \\ &= 2R_{-1}(M_U) + 2\sum_{k<l} \mu_k\mu_l \\ &= 0.\end{aligned}$$

Hence, $\sum_{k<l} \mu_k\mu_l = -R_{-1}(M_U)$.

From the definition of the Hermitian-Randić energy of a mixed graph, we have

$$\begin{aligned}\mathcal{E}_{R_H}^2(M) &= \left(\sum_{k=1}^{n} |\mu_k|\right)^2 \\ &= \sum_{k=1}^{n} \mu_k^2 + \sum_{1\le k\ne l\le n} |\mu_k\mu_l| \\ &= 2R_{-1}(M_U) + 2\sum_{k<l} |\mu_k\mu_l| \\ &\ge 2R_{-1}(M_U) + 2|\sum_{k<l} \mu_k\mu_l| \\ &= 4R_{-1}(M_U).\end{aligned}$$

Combining with Lemma 3.7, we have

$$\mathcal{E}_{R_H}(M) \ge 2\sqrt{R_{-1}(M_U)} \ge 2\sqrt{\frac{n}{2(n-1)}} = \sqrt{\frac{2n}{n-1}}.$$

So,

$$\sqrt{\frac{2n}{n-1}} \le \mathcal{E}_{R_H}(M) \le n.$$

From the proof above and Lemma 3.7, we know that the equality in the lower bound holds if and only if M_U is a complete graph and $\mu_k\mu_l \ge 0$ or $\mu_k\mu_l \le 0$, for all $1 \le k < l \le n$. Note that $\sum_{k=1}^{n} \mu_k = 0$ and M has no isolated vertices, so the former case can not happen. Hence, the equality in the lower bound holds if and only if M_U is a complete graph and $\mu_1 = -\mu_n \ne 0$, $\mu_j = 0$, $j = 2, \ldots, n-1$.

This completes the proof. ■

Remark 3.9. It should be pointed out that when M is a complete mixed graph, its Hermitian-Randić spectrum is not unique. For example, let $M_U = K_3$, if all edges of $E(M)$ are oriented, then we have $\mu_1 = -\mu_3 = \frac{\sqrt{3}}{2}$, $\mu_2 = 0$, then we can obtain the lower bound in Theorem 3.8.

If there exist some edges of $E(M)$ are undirected, then we can not obtain the lower bound in Theorem 3.8. For example, if $(r_h)_{12} = (r_h)_{32} = \frac{i}{2}$, $(r_h)_{13} = \frac{1}{2}$, then $\mu_1 = 1$ and $\mu_2 = \mu_3 = -\frac{1}{2}$. Hence, the problem of determining all complete mixed graphs for which the lower bound in Theorem 3.8 is attained appears to be somewhat more difficult.

To deduce more bounds on $\mathcal{E}_{R_H}(M)$, the following lemma is needed.

Lemma 3.10. *[8] Let $\boldsymbol{x},\ \boldsymbol{y} \in \mathbb{R}^n$ and let $A(\boldsymbol{x}) = \frac{1}{n}\sum_{i=1}^n x_i$, $A(\boldsymbol{y}) = \frac{1}{n}\sum_{i=1}^n y_i$. If $\phi \le x_i \le \Phi$ and $\gamma \le y_i \le \Gamma$, then*

$$\left|\frac{1}{n}\sum_{i=1}^n x_i y_i - \frac{1}{n^2}\sum_{i=1}^n x_i \sum_{i=1}^n y_i\right| \le \sqrt{(\Phi - A(x))(A(x)-\phi)(\Gamma - A(y))(A(y)-\gamma)}.$$

Now we turn to new bounds on $\mathcal{E}_{R_H}(M)$.

Theorem 3.11. *Let M be a mixed graph of order n and M_U be its underlying graph. Let $\mu_1 \ge \mu_2 \ge \cdots \ge \mu_n$ be the Hermitian-Randić spectrum of $R_H(M)$. Then*

$$\mathcal{E}_{R_H}(M) \ge \frac{2R_{-1}(M_U) + n\alpha\beta}{\alpha+\beta}, \tag{1}$$

where $\alpha = \min\limits_{1\le i\le n}\{|\mu_i|\}$, $\beta = \max\{\mu_1, |\mu_n|\}$.

Proof. Note that

$$\begin{aligned}\mathcal{E}^2_{R_H}(M) &= \left(\sum_{j=1}^n |\mu_j|\right)^2 = \sum_{j=1}^n |\mu_j|^2 + \sum_{1\le i\ne j\le n} |\mu_i||\mu_j| \\ &= 2R_{-1}(M_U) + \sum_{1\le i\ne j\le n} |\mu_i||\mu_j|.\end{aligned} \tag{2}$$

Let $S = \sum_{1\le i\ne j\le n} |\mu_i||\mu_j|$, $x_i = |\mu_i|$ and $y_i = \mathcal{E}_{R_H}(M) - |\mu_i|$, $i = 1, 2, \ldots, n$. Then $S = \sum_{i=1}^n x_i y_i$.

From the definitions of α and β, we have $\alpha \le x_i \le \beta$ and $\mathcal{E}_{R_H}(M) - \beta \le y_i \le \mathcal{E}_{R_H}(M) - \alpha$. In addition, let $A(\mathbf{x}) = \frac{1}{n}\sum_{i=1}^n x_i = \frac{\mathcal{E}_{R_H}(M)}{n}$ and $A(\mathbf{y}) = \frac{1}{n}\sum_{i=1}^n y_i = \frac{(n-1)\mathcal{E}_{R_H}(M)}{n}$. Hence by Lemma 3.10, we have

$$\begin{aligned}\left|\frac{S}{n} - \frac{(n-1)\mathcal{E}^2_{R_H}(M)}{n^2}\right| &\le \sqrt{\left(\beta - \frac{\mathcal{E}_{R_H}(M)}{n}\right)\left(\frac{\mathcal{E}_{R_H}(M)}{n} - \alpha\right)} \\ &\quad \cdot \sqrt{\left[\mathcal{E}_{R_H}(M) - \alpha - \frac{(n-1)\mathcal{E}_{R_H}(M)}{n}\right]} \\ &\quad \cdot \sqrt{\left[\frac{(n-1)\mathcal{E}_{R_H}(M)}{n} - (\mathcal{E}_{R_H}(M) - \beta)\right]} \\ &= \sqrt{\left(\beta - \frac{\mathcal{E}_{R_H}(M)}{n}\right)^2\left(\frac{\mathcal{E}_{R_H}(M)}{n} - \alpha\right)^2}.\end{aligned}$$

It follows that

$$S \ge \mathcal{E}^2_{R_H}(M) + n\alpha\beta - (\alpha+\beta)\mathcal{E}_{R_H}(M).$$

This together with (2) implies that

$$\mathcal{E}^2_{R_H}(M) = 2R_{-1}(M_U) + S \geq 2R_{-1}(M_U) + \mathcal{E}^2_{R_H}(M) + n\alpha\beta - (\alpha+\beta)\mathcal{E}_{R_H}(M).$$

So,

$$\mathcal{E}_{R_H}(M) \geq \frac{2R_{-1}(M_U) + n\alpha\beta}{\alpha+\beta}.$$

This completes the proof. ■

Note that the right-hand side of (1) is a non-decreasing function on $\alpha \geq 0$. Combining with Theorem 3.4, we have the following corollary.

Corollary 3.12. *Let M be a mixed graph of order n and M_U be its underlying graph. Let $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_n$ be the Hermitian-Randić spectrum of $R_H(M)$. Then*

$$\mathcal{E}_{R_H}(M) \geq \frac{2R_{-1}(M_U)}{\beta},$$

where $\beta = \max\{\mu_1, |\mu_n|\}$. The equality holds if and only if $R^2_H(M) = c\mathbf{I}_n$, where c is a constant such that $|\mu_i|^2 = c$ for all i.

In particular, if M is a connected mixed bipartite graph, then we have the following theorem.

Theorem 3.13. *Let M be a connected mixed bipartite graph of order n and M_U be its underlying graph. Let $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_n$ be the Hermitian-Randić spectrum of $R_H(M)$. Then*

$$\mathcal{E}_{R_H}(M) \geq 2\left(\frac{R_{-1}(M_U) + \lfloor \frac{n}{2} \rfloor \alpha\mu_1}{\alpha+\mu_1}\right), \tag{3}$$

where $\alpha = \min\limits_{1\leq i\leq \lfloor \frac{n}{2} \rfloor}\{|\mu_i|\}$.

Proof. Note that M_U is a bipartite graph. By Corollary 2.3 (3), we have $\mu_i = -\mu_{n+1-i}$ and $\mu_i \geq 0$ for $i = 1, 2, \ldots, \lfloor \frac{n}{2} \rfloor$. Therefore,

$$\begin{aligned}\mathcal{E}^2_{R_H}(M) &= \left(2\textstyle\sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \mu_i\right)^2 = 4(\textstyle\sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \mu_i^2 + \sum_{1\leq i\neq j\leq \lfloor \frac{n}{2} \rfloor} \mu_i\mu_j) \\ &= 4R_{-1}(M_U) + 4\textstyle\sum_{1\leq i\neq j\leq \lfloor \frac{n}{2} \rfloor} \mu_i\mu_j.\end{aligned} \tag{4}$$

Let $T = \sum_{1\leq i\neq j\leq \lfloor \frac{n}{2} \rfloor} \mu_i\mu_j$, $x_i = \mu_i$ and $y_i = \frac{\mathcal{E}_{R_H}(M)}{2} - \mu_i$, $i = 1, 2, \ldots, \lfloor \frac{n}{2} \rfloor$. Then $T = \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} x_i y_i$.

From the definition of α, we have $\alpha \leq x_i \leq \mu_1$ and $\frac{\mathcal{E}_{R_H}(M)}{2} - \mu_1 \leq y_i \leq \frac{\mathcal{E}_{R_H}(M)}{2} - \alpha$. In addition, let $A(\mathbf{x}) = \frac{1}{\lfloor \frac{n}{2} \rfloor}\sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} x_i = \frac{\mathcal{E}_{R_H}(M)}{2\lfloor \frac{n}{2} \rfloor}$ and $A(\mathbf{y}) = \frac{1}{\lfloor \frac{n}{2} \rfloor}\sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} y_i = \frac{(\lfloor \frac{n}{2} \rfloor - 1)\mathcal{E}_{HR}(M)}{2\lfloor \frac{n}{2} \rfloor}$. Hence by Lemma 3.10, we have

$$\left|\frac{T}{\lfloor \frac{n}{2} \rfloor} - \frac{(\lfloor \frac{n}{2} \rfloor - 1)\mathcal{E}^2_{R_H}(M)}{4\lfloor \frac{n}{2} \rfloor^2}\right| \leq \sqrt{\left(\mu_1 - \frac{\mathcal{E}_{R_H}(M)}{2\lfloor \frac{n}{2} \rfloor}\right)^2\left(\frac{\mathcal{E}_{R_H}(M)}{2\lfloor \frac{n}{2} \rfloor} - \alpha\right)^2}.$$

It follows that

$$T \geq \frac{\mathcal{E}^2_{R_H}(M)}{4} + \lfloor \frac{n}{2} \rfloor \alpha\mu_1 - (\alpha+\mu_1)\frac{\mathcal{E}_{R_H}(M)}{2}.$$

This together with (4) implies that

$$\mathcal{E}^2_{R_H}(M) = 4R_{-1}(M_U) + 4T \geq 4R_{-1}(M_U) + \mathcal{E}^2_{R_H}(M) + 4\lfloor\frac{n}{2}\rfloor\alpha\mu_1 - 2(\alpha+\mu_1)\mathcal{E}_{R_H}(M).$$

So,

$$\mathcal{E}_{R_H}(M) \geq 2\left(\frac{R_{-1}(M_U) + \lfloor\frac{n}{2}\rfloor\alpha\mu_1}{\alpha+\mu_1}\right).$$

This completes the proof. ■

Note that the right-hand side of (3) is a non-decreasing function on $\alpha \geq 0$. Combining with Theorem 3.4, we have the following corollary.

Corollary 3.14. *Let M be a connected mixed bipartite graph of order n and M_U be its underlying graph. Let $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_n$ be the Hermitian-Randić spectrum of $R_H(M)$. Then*

$$\mathcal{E}_{R_H}(M) \geq \frac{2R_{-1}(M_U)}{\mu_1},$$

the equality holds if and only if $R^2_H(M) = c\mathbf{I}_n$, where c is a constant such taht $|\mu_i|^2 = c$ for all i.

4 Hermitian-Randić energy of trees

In [1], the authors proved that the skew energy of a directed tree is independent of its orientation. In [11], the authors showed that the skew Randić energy of a directed tree has the same result. In this section, we will show that the Hermitian-Randić energy also has the same result. In the beginning of this section, we first to characterize the mixed graphs with cut-edge.

Theorem 4.1. *Let M be a mixed graph of order n and $e = uv$ is an edge of M. If uv is a cut-edge of M_U, where M_U is the underlying graph of M. Then the spectrum and energy of $R_H(M)$ are unchanged when the edge uv is replaced with a single arc uv or vu and vice versa.*

Proof. Let $e = uv$ be a cut-edge of M_U. Suppose M_1 is the graph obtained from M by replacing the edge uv with the arc uv or vu. Let M' and M'_1 be real elementary subgraph of order k of M and M_1, respectively. If M' do not contain the cut-edge uv, then M' is also a real elementary subgraph of M_1, that is, $M' = M'_1$. So we have

$$(-1)^{r(M')+l(M')}2^{s(M')}\prod_{v_i\in V(M')}\frac{1}{d_{M_U}(v_i)} = (-1)^{r(M'_1)+l(M'_1)}2^{s(M'_1)}\prod_{v_i\in V(M'_1)}\frac{1}{d_{M_U}(v_i)}. \quad (5)$$

If M' contains the cut-edge uv, then there is a real elementary subgraph M'_1 of M_1 only differ from M' on uv. Since uv is a cut-edge of M_U, uv is not contained in any cycles of M_U. Hence we have

$$(-1)^{r(M')+l(M')}2^{s(M')}\prod_{v_i\in V(M')}\frac{1}{d_{M_U}(v_i)} = (-1)^{r(M'_1)+l(M'_1)}2^{s(M'_1)}\prod_{v_i\in V(M'_1)}\frac{1}{d_{M_U}(v_i)}. \quad (6)$$

Combining with (5) and (6), we have

$$\begin{aligned}
&a_k(M) - a_k(M_1) \\
&= \sum_{M'} (-1)^{r(M')+l(M')} 2^{s(M')} \prod_{v_i \in V(M')} \frac{1}{d_{M_U}(v_i)} \\
&- \sum_{M_1'} (-1)^{r(M_1')+l(M_1')} 2^{s(M_1')} \prod_{v_i \in V(M_1')} \frac{1}{d_{M_U}(v_i)} \\
&= 0,
\end{aligned}$$

for any integer k. Thus $Sp_{R_H}(M) = Sp_{R_H}(M_1)$. Moreover, $\mathcal{E}_{R_H}(M) = \mathcal{E}_{R_H}(M_1)$.

Similarly, we can prove that $Sp_{R_H}(M) = Sp_{R_H}(M_2)$ and $\mathcal{E}_{R_H}(M) = \mathcal{E}_{R_H}(M_2)$, where M_2 is the mixed graph obtained from M by replacing the arc uv or vu with the edge uv. ■

Thus, the Hermitian-Randić spectrum and Hermitian-Randić energy are invariants when reversing the cut-arc's orientation or unorienting it or orienting an undirected cut-edge. By applying Theorem 4.1, we can obtain the following corollaries.

Corollary 4.2. *Let T be a mixed tree of order n, and T' be the mixed tree obtained from T by reversing the orientations of all the arcs incident with a particular vertex of T. Then $\mathcal{E}_{R_H}(T) = \mathcal{E}_{R_H}(T')$.*

Corollary 4.3. *Let T be a mixed tree and T_U be its underlying graph. Then*

(1) The Hermitian-Randić energy of T is independent of its orientation of the arc set.

(2) The Hermitian-Randić energy of T is the same as the Randić energy of T_U.

References

[1] C. Adiga, R. Balakrishnan, W. So, The skew energy of a digraph, *Linear Algebra Appl.* **432** (2010) 1825–1835.

[2] J. A. Bondy, U. S. R. Murty, *Graph Theory with Applications*, Elsevier, New York, 1976.

[3] Ş. B. Bozkurt, D. Bozkurt, Randić energy and Randić Estrada index of a graph, *Eur. J. Pure Appl. Math.* **5** (2012) 88–96.

[4] Ş. B. Bozkurt, D. Bozkurt, Sharp upper bounds for energy and Randić energy, *MATCH Commun. Math. Comput. Chem.* **70** (2013) 669–680.

[5] Ş. B. Bozkurt, A. D. Güngör, I. Gutman, Randić spectral radius and Randić energy, *MATCH Commun. Math. Comput. Chem.* **64** (2010) 321–334.

[6] Ş. B. Bozkurt, A. D. Güngör, I. Gutman, A. S. Çevik, Randić matrix and Randić energy, *MATCH Commun. Math. Comput. Chem.* **64** (2010) 239–250.

[7] Xiaolin Chen, Xueliang Li, Yingying Zhang, 3-Regular mixed graphs with optimum Hermitian energy, *Linear Algebra Appl.* **496** (2016) 475–486.

[8] S. S. Dragomir, A generaization of Grüss's inequaity in inner product spaces and applications, *J. Math. Anal. Appl.* **237** (1999) 74–82.

[9] Shicai Gong, Guanghui Xu, The characteristic polynomial and the matchings polynomial of a weighted oriented graph, *Linear Algebra Appl.* **436** (2012) 3597–3607.

[10] Ran Gu, Fei Huang, Xueliang Li, General Randić matrix and general Randić energy, *Trans. Comb.* **3** no. 3 (2014) 21–33.

[11] Ran Gu, Fei Huang, Xueliang Li, Skew Randić matrix and skew Randić energy, *Trans. Comb.* **5** no. 1 (2016) 1–14.

[12] Ran Gu, Xueliang Li, Jinfeng Liu, Note on three results on Randić energy and incidence energy, *MATCH Commun. Math. Comput. Chem.* **73** (2015) 61–71.

[13] K. Guo, B. Mohar, Hermitian adjacency matrix of digraphs and mixed graphs, *J. Graph Theory* **00** (2016) 1–32.

[14] I. Gutman, B. Furtula, Ş. B. Bozkurt, On Randić energy, *Linear Algebra Appl.* **422** (2014) 50–57.

[15] I. Gutman, Xueliang Li, Jianbin Zhang, Graph energy, in: M. Dehmer, F. Emmert-Streib (Eds.), *Analysis of Complex Network: From Biology to Linguistics*, Wiley-VCH Verlag, Weinheim, 2009, pp. 145–174.

[16] Yaoping Hou, Tiangang Lei, Characteristic polynomials of skew-adjacency matrices of oriented graphs, *Electron. J. Combin.* **18** (2011) P#156.

[17] Jianxi Li, Jiming Guo, Wai Chee Shiu, A Note on Randić energy, *MATCH Commun. Math. Comput. Chem.* **74** (2015) 389–398.

[18] Xueliang Li, I. Gutman, *Mathematical Aspects of Randic-type Molecular Structure Descriptors*, Univ. Kragujevac, Kragujevac, 2006.

[19] Xueliang Li, Huishu Lian, Skew energy of oriented graphs, Chapter 8 in "I. Gutman, X.L. Li (Eds.), Energies of Graphs-Theory and Applications", Mathematical Chemistry Monograph No.17, 2016, pp.191–236. Or available at https://arxiv.org/abs/1304.5707, 2013.

[20] Xueliang Li, Yongtang Shi, A survey on the Randic index, *MATCH Commun. Math. Comput. Chem.* **59** (2008) 127–156.

[21] Xueliang Li, Yongtang Shi, I. Gutman, *Graph Energy*, Springer, New York, 2012.

[22] Xueliang Li, Jianfeng Wang, Randić energy and Randić eigenvalues, *MATCH Commun. Math. Comput. Chem.* **73** (2015) 73–80.

[23] Xueliang Li, Yiting Yang, Best lower and upper bounds for the Randić index R_{-1} of chemical trees, *MATCH Commun. Math. Comput. Chem.* **52** (2004) 147–156.

[24] Xueliang Li, Yiting Yang, Sharp bounds for the general Randić index, *MATCH Commun. Math. Comput. Chem.* **51** (2004) 155–166.

[25] Jianxi Liu, Xueliang Li, Hermitian-adjacency matrices and Hermitian energies of mixed graphs, *Linear Algebra Appl.* **466** (2015) 182–207.

[26] B. Mohar, Hermitian adjacency spectrum and switching equivalence of mixed graphs, *Linear Algebra Appl.* **489** (2016) 324–340.

[27] M. Randić, On characterization of molecular branching, *J. Amer. Chem. Soc.* **97** (1975) 6609–6615.

[28] M. Randić, On history of the Randić index and emerging hostility toward chemical graph theory, *MATCH Commun. Math. Comput. Chem.* **59** (2008) 5–124.

[29] Yongtang Shi, Note on two generalizations of the Randić index, *Appl. Math. Comput.* **265** (2015) 1019–1025.

[30] Guihai Yu, Xin Liu, Hui Qu, Singularity of Hermitian (quasi-)Laplacian matrix of mixed graphs, *Appl. Math. Comput.* **293** (2017) 287–292.

[31] Guihai Yu, Hui Qu, Hermitian Laplacian matrix and positive of mixed graphs, *Appl. Math. Comput.* **269** (2015) 70–76.